


Restoration of the Lippe floodplain and the burbot (*Lota lota*)

Margret Bunzel-Drücke,
Matthias Scharf & Olaf Zimball




Arbeitsgemeinschaft Biologischer
Umweltschutz im Kreis Soest e.V.
Bad Sassendorf-Lohne

Distribution of the burbot in NRW in the 19. century



Distribution of the burbot in NRW in the 19. century



Distribution of the burbot in NRW
in 2000 (stocking excluded)

Regulated river:

**stone riprap often used
by adult burbot**



Characteristics of spawning grounds

- cold (1 - 4 °C)
- shallow
- rich in oxygen
- gentle current at best
- low predation pressure



Lippe burbot spawn in small tributaries reaching the floodplain.


Burbot eggs

diameter ca. 1 mm



Characteristics of burbot larval habitats

- no (strong) current
- shallow areas
- rich in zooplankton
- low predation pressure and not much competition



length ca. 4 mm

Where are such habitats in spring?

Müller (1960): Biologie der Quappe zwischen Elbe und Oder (Zeitschrift für Fischerei 9 N.F.)


Foto: Spree on 06.04.1954




Inundated floodplain



Burbot habitats in the Lippe floodplain: near natural conditions




Shortened river course



incision (down-cutting) of the stream channel



Lippe floodplain: loss of flood channels





Floodplain still flooded sometimes ...

... but ...
**summer
embankments
separate river
from floodplain.**



Restoration of the Lippe floodplain: project partners



The State of North-Rhine/Westphalia



Lippeverband



The City of Hamm LIFE projects



The District of Soest



Nordrhein-Westfalen-Stiftung
Naturschutz, Heimat- und Kulturpflege



Arbeitsgemeinschaft Biologischer
Umweltschutz im Kreis Soest e.V.





**Restoration measures:
river bed is widened
and raised up**



Before ...

... and after


the restoration.



- river bed widened, lengthened and raised up
- in the floodplain lakes and flood channels restored




Extent of Lippe restoration measures




Study method: standardized electrofishing



Reaction of the Lippe burbot population






5 km

Number of young-of-the-year burbot 2000



restored floodplain



no record



1 – 5 individuals




6 – 10 individuals

11 – 50 individuals


Restoration of the Lippe floodplain and the burbot


Margret Bunzel-Drücke, Matthias Scharf & Olaf Zimball, 24.09.2015

Reaction of the Lippe burbot population




Reaction of the Lippe burbot population





Numbers of adult and 0+ burbot 2013

restored floodplain



no record

1 – 5 Individuals

6 – 10 Individuals

11 – 50 Individuals

> 50 Individuals





Thank you for your attention!

